Eastern Health Laboratory Medicine	Section: Management System\Eastern Hea	alth\Client Services\Collection Manual\
	Title: Collecting a 5-HIAA Random	Urine Number: 14043
	Specimen	Version: 2.0
	Effective Date: 2/27/2019 Sta	atus: Current

How to Collect a 5-HIAA Random Urine Specimen

Please read all this material **before** you start your urine collection.

Samples not labelled correctly and/or do not have a complete requisition will not be processed. To ensure correct test results, follow instructions carefully.

Before Collecting the Random Urine Specimen:

Certain foods and medications can interfere with the test. The following a diet and medication restrictions must be followed for 72 hours before and during all of the urine collection (24 hours):

Avoid all of the following foods and compounds.

Coffee (Caffeine) Nicotine (smoking) Banana Chocolate Eggplant Fruit and Juice Pineapple Cola drink Seafoods

Pumpkins Soy products Seaweed Spinach Liver Melon family Date Kiwi Papaya

Grapefruit Prune and Plum Plantain Tomato and tomato-based products Теа Avocado Cantaloupe Nuts (Walnuts, almonds, pecans, pistachios, etc.)

Avoid all of the following over-the-counter medications.

ASA (e.g. Aspirin) Acetaminophen (e.g. Tylenol) Cough, Cold and Flu medications Antihistamines

Ask your doctor whether it is safe for you to stop taking the following prescription medications as they may affect the test results.

Anti-hypertensive drugs Levodopa (L-dopa) Chloral hydrate Dopamine

Erythromycin Niacin Phenothiazine Quinidine

Quinine Riboflavin Tetracvcline MAO inhibitors

How to collect the samples:

- Collect the random urine in the clean sterile container provided to you.
- Bring the sample to your local laboratory within 3 hours of collection.